AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

<u>Drug Requested</u> : (Select drug below)				
□ deferiprone (Ferriprox®) tablets	□ Ferriprox [®] (deferiprone) solution	☐ Ferriprox® 2-day (deferiprone) tablets		
MEMBER & PRESCRIBE	R INFORMATION: Authorizat	ion may be delayed if incomplete.		
Member Name:				
Member AvMed #:		Date of Birth:		
Prescriber Name:				
Prescriber Signature:		Date:		
Office Contact Name:				
	one Number: Fax Number:			
DEA OR NPI #:				
DRUG INFORMATION: A	authorization may be delayed if incon	nplete.		
Drug Form/Strength:				
	sing Schedule: Length of Therapy:			
Diagnosis:	agnosis: ICD Code, if applicable:			
Weight:	Date:			
Quantity Limits: Maximum 99	mg/kg/day (actual body weight) in t	wo divided doses		
support each line checked, all docu	neck below all that apply. All criteriumentation, including lab results, diagnetic check the diagnosis below that apply.	gnostics, and/or chart notes, must be		
Diagnosis: Transfusiona	l hemosiderosis due to thalass	omia syndromo		
		emia syndrome		
Initial Authorization: 6 mor				
☐ Member is 3 years of age or				
	ransfusional hemosiderosis due to the blood cells, approximately 20 units for	alassemia syndrome (i.e., transfusion of or a 40 kg patient)		

(Continued on next page)

Member's serum ferritin levels are consistently >1,000 mcg/L (submit serum ferritin labs done within the last 30 days)
Member's current weight:
Member is currently established on <u>ONE</u> of the following chelator therapies for at least 90 consecutive days (verified by pharmacy paid claims):
□ deferoxamine (Desferal)
□ deferasirox (Exjade, Jadenu)
Member continues to have ONE of the following after three (3) months of chelator therapy:
□ Serum ferritin in excess of 2,500 mcg/L
□ Liver iron concentration is >7 mg Fe/g dry weight (submit liver biopsy, MRI or other FDA-approved test results)
□ Cardiac magnetic resonance imaging (MRI T2*) is \leq 10 ms (submit MRI T2* lab results)
Baseline absolute neutrophil count (ANC) is $> 1.5 \times 10^9/L$ and ANC will continue to be monitored weekly while on therapy (submit current labs)
If requesting brand Ferriprox, documentation of trial and intolerable life-endangering adverse event with generic deferiprone must be submitted
Ferriprox solution may be approved for members aged 3-10 years only. If requesting Ferriprox solution for members ≥ 11 years of age, documentation that member is unable to ingest any solid dosage form
must be submitted
· · · · · · · · · · · · · · · · · · ·
must be submitted
must be submitted Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome
must be submitted Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months.
must be submitted Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is $> 1.5 \times 10^9$ /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other
must be submitted Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is $> 1.5 \times 10^9$ /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results)
Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is $> 1.5 \times 10^9$ /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results) Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L Serum ferritin has decreased by $\geq 20\%$ from baseline or has been maintained at a level that is $\geq 20\%$
Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is $> 1.5 \times 10^9$ /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results) Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L Serum ferritin has decreased by $\ge 20\%$ from baseline or has been maintained at a level that is $\ge 20\%$ below baseline level (submit current serum ferritin labs) Diagnosis: Transfusional iron overload in members with sickle cell disease or other
Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is $> 1.5 \times 10^9$ /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results) Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L Serum ferritin has decreased by $\ge 20\%$ from baseline or has been maintained at a level that is $\ge 20\%$ below baseline level (submit current serum ferritin labs) Diagnosis: Transfusional iron overload in members with sickle cell disease or other memias
Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is $> 1.5 \times 10^9$ /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results) Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L Serum ferritin has decreased by $\ge 20\%$ from baseline or has been maintained at a level that is $\ge 20\%$ below baseline level (submit current serum ferritin labs) Diagnosis: Transfusional iron overload in members with sickle cell disease or other memias ial Authorization: 12 months
Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome uthorization: 12 months. Member's ANC is > 1.5×10^9 /L (submit current lab results) Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results) Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L Serum ferritin has decreased by $\geq 20\%$ from baseline or has been maintained at a level that is $\geq 20\%$ below baseline level (submit current serum ferritin labs) Diagnosis: Transfusional iron overload in members with sickle cell disease or other memias ial Authorization: 12 months Member is 3 years of age or older Member has a diagnosis of transfusional iron overload associated with sickle cell disease or other

(Continued on next page)

(1	Baseline absolute neutrophil count (ANC) is $> 1.5 \times 10^9/L$ and ANC will continue to be monitored weekly while on therapy (submit current labs)
		iagnosis: Transfusional iron overload in members with sickle cell disease or other nemias
		<u>ithorization</u> : 12 months.
Į	3	Liver iron concentration has decreased by at least 4 mg of Fe/g dry weight from baseline or has been maintained at a level that is at least 4 mg of Fe/g dry weight below baseline level since last approval (submit current MRI results)
[ב	Member's ANC is $> 1.5 \times 10^9/L$ (submit current lab results)
Į	ב	Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L

Medication being provided by a Specialty Pharmacy - PropriumRx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

**Use of samples to initiate therapy does not meet step edit/preauthorization criteria. **

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.