

# AvMed

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization may be delayed.**

**Drug Requested:** Firdapse<sup>®</sup> (amifampridine phosphate)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member AvMed #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

### **Recommended Dosage:**

Age and body weight	Initial daily dosage	Titration regimen	Maximum single dose	Maximum total daily maintenance dosage
<ul style="list-style-type: none"><li>Adults (any weight)</li><li>Pediatric patients weighing 45 kg or more</li></ul>	15 mg to 30 mg daily, in 3 to 4 divided doses	Increase total daily dosage by 5 mg every 3 or 4 days	20 mg	80 mg given in divided doses
<ul style="list-style-type: none"><li>Pediatric patients weighing less than 45 kg</li></ul>	5 mg to 15 mg daily, in 3 to 4 divided doses	Increase total daily dosage by 2.5 mg every 3 or 4 days	10 mg	40 mg given in divided doses

**Quantity Limit:** 240 tablets per 30 days

(Continued on next page)

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Initial Authorization: 6 months**

- Medication must be prescribed by or in consultation with a neurologist
- Member must be 6 years of age or older
- Member must have a diagnosis of Lambert-Eaton myasthenic syndrome
- Diagnosis has been confirmed by **ONE** of the following (**must submit labs for documentation**):
  - Presence of anti-P/Q-type voltage-gated calcium channel (VGCC) antibodies
  - A confirmatory electrodiagnostic study [e.g., repetitive nerve stimulation (RNS), needle electromyography (EMG), single-fiber electromyography (SFEMG)]
- Must submit chart notes of moderate to severe muscle weakness that interferes with function
- Attestation that other differential diagnoses such as Myasthenia gravis have been ruled out
- Attestation that the member does **NOT** have a history of seizures or take medications that lower the seizure threshold (e.g., bupropion, tramadol, amphetamines, theophylline)
- Provider attests the member is **NOT** using alcohol
- Member is **NOT** receiving Firdapse® in combination with similar potassium channel blockers, such as Ampyra® (dalfampridine), or used in combination with compounded formulation of 3,4 diaminopyridine
- Provider must submit a baseline assessment or chart notes documenting **ONE** of the following measures:
  - Dynamometry
  - Timed 25 Foot Walk test
  - Timed Up and Go (TUG) test

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Must submit chart notes of a positive clinical symptomatic response to Firdapse® therapy with improvement from the initial submitted baseline assessment (**current assessment must be submitted**):
  - Dynamometry
  - Timed 25 Foot Walk test (a quantitative mobility and leg function performance test based on a timed 25-foot walk; an average increase of more than 20% in the timed 25-foot walk may indicate a significant change in gait)
  - Timed Up and Go (TUG) test (assesses patient's function, weakness and mobility. The test measures the time it takes for patients to rise from a chair, walk a short distance, return to their chair and climb stairs approximately three times; >30% time increase from baseline indicates deterioration)
    - 11–20 seconds is within normal limits for frail elderly and disabled patients
    - Greater than 20 seconds suggests the person needs assistance and indicates further examination and intervention may be required
    - 30 seconds or more suggests that the person may be prone to falls

(Continued on next page)

Medication being provided by Specialty Pharmacy - PropriumRx

*Not all drugs may be covered under every Plan.*

*If a drug is non-formulary on a Plan, documentation of medical necessity will be required.*

*\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\**

*\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\**