

AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Trikafta[®] (elexacaftor/tezacaftor/ivacaftor and ivacaftor)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

Maximum Approved Dose:

- **Children ≥ 2 years to < 6 years weighing < 14 kg:** Oral: 1 packet (containing elexacaftor 80 mg/tezacaftor 40 mg/ivacaftor 60 mg) in the morning and 1 packet (containing ivacaftor 59.5 mg) in the evening
- **Children ≥ 2 years to < 6 years weighing > 14 kg:** Oral: 1 packet (containing elexacaftor 100 mg/tezacaftor 50 mg/ivacaftor 75 mg) in the morning and 1 packet (containing ivacaftor 75 mg) in the evening
- **Children ≥ 6 years to < 12 years weighing < 30 kg:** Oral: 2 tablets (each containing elexacaftor 50 mg/tezacaftor 25 mg/ivacaftor 37.5 mg) in the morning and ivacaftor 75 mg in the evening, approximately 12 hours apart
- **Children ≥ 6 years to < 12 years weighing ≥ 30 kg:** Oral: 2 tablets (each containing elexacaftor 100 mg/tezacaftor 50 mg/ivacaftor 75 mg) in the morning and ivacaftor 150 mg in the evening, approximately 12 hours apart
- **Children ≥ 12 years, Adolescents and Adults:** Oral: 2 tablets (each containing elexacaftor 100 mg/tezacaftor 50 mg/ivacaftor 75 mg) in the morning and one ivacaftor 150 mg tablet in the evening, approximately 12 hours apart

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- Member is **2 years of age or older** with a diagnosis of Cystic Fibrosis
- Member has **at least one** of the F508 del mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or a mutation in the CFTR gene as confirmed by an FDA-cleared test that is responsive to elexacaftor/tezacaftor/ivacaftor (**Test results must be attached**)
- Prescribing physician is a pulmonologist or has consulted with a pulmonologist who specializes in the treatment of Cystic Fibrosis
- Baseline FEV1 within the last 30 days must be submitted, unless the member is unable to perform a pulmonary function test (**Test results must be attached**)
- Provider attests that baseline ophthalmic examination to monitor lens opacities/cataracts has been completed for pediatric members
- Baseline liver function tests have been completed prior to initiating therapy and will be completed annually (**Labs must be attached**)
- Member does **NOT** have severe hepatic impairment (Child-Pugh Class C)
- Number of pulmonary exacerbations or hospitalizations in the preceding 6 months must be noted: _____
- Baseline body mass index must be noted: _____
- Member will **NOT** be taking Trikafta® in combination with any other CFTR modulator therapy (i.e., Symdeko®, Orkambi®, Kalydeco®) **NOTE:** Concurrent therapy with these agents will **NOT** be approved
- Member will avoid concomitant use of strong CYP3A inducers (e.g., rifampin, carbamazepine, phenytoin, phenobarbital, St. John's wort) and strong or moderate CYP3A inhibitors (e.g., fluconazole, itraconazole)

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member continues to meet initial criteria
- Member has demonstrated disease response as indicated by **one or more** of the following (**must submit current labs and chart notes**):
 - Decreased pulmonary exacerbations or hospitalizations compared to pretreatment baseline
 - Stabilization of lung function as measured by FEV1 within the last year compared to baseline
 - Improvement in quality of life, weight gain, or growth
- Member has **NOT** received a lung transplant
- Member has experienced an absence of unacceptable toxicity from therapy (i.e. elevated transaminases (ALT or AST), development of cataracts or lens opacities)

Date of initiation of Trikafta [®] therapy: _____	Reauthorization Date: _____
Baseline FEV1 (last FEV1 prior to starting Trikafta [®]): _____	Current FEV1 (FEV1 AFTER last dose of Trikafta [®]): _____
Baseline Weight: _____	Current weight: _____
Baseline BMI: _____	Current BMI: _____
Number of hospitalizations since last approval of Trikafta [®] must be noted _____	

Medication being provided by a Specialty Pharmacy – Proprium Rx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.