

Table 1

COVID-19 vaccination recommendations have changed. Find the latest recommendations at www.cdc.gov/covidschedule
Recommended Adult Immunization Schedule for ages 19 years or older, United States, 2023

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
COVID-19	2- or 3- dose primary series and booster (See Notes)			
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually			
Influenza live, attenuated (LAIV4)	1 dose annually			
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			For healthcare personnel, see notes
Varicella (VAR)	2 doses (if born in 1980 or later)	2 doses		
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (see notes)		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (see notes)			See Notes
				See Notes
Hepatitis A (HepA)	2, 3, or 4 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations			
	19 through 23 years			
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			

 Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection
 Recommended vaccination for adults with an additional risk factor or another indication
 Recommended vaccination based on shared clinical decision-making
 No recommendation/ Not applicable