

AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Kerendia[®] (finerenone)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

Quantity Limit: 30 tablets per 30 days

Kerendia[®] Initial Dosing Recommendations:

eGFR (mL/min/1.73m ²)	Starting Dose
≥ 60	20 mg once daily
≥ 25 to < 60	10 mg once daily
< 25	Not Recommended

(Continued on next page)

Dose Adjustment Based on Current Serum Potassium Concentration and Current Kerendia® Dose			
		10 mg once daily	20 mg once daily
Current Serum Potassium (mEq/L)	≤ 4.8	Increase the dose to 20 mg once daily*	Maintain 20 mg once daily
	> 4.8 – 5.5	Maintain 10 mg once daily	Maintain 20 mg once daily
	> 5.5	Withhold Kerendia®. Consider restarting at 10 mg once daily when serum potassium ≤ 5.0 mEq/L	Withhold Kerendia®. Restart at 10 mg once daily when serum potassium ≤ 5.0 mEq/L

*If eGFR has decreased by more than 30% compared to previous measurement, maintain 10 mg dose.

CLINICAL CRITERIA: Check below all that apply. **All criteria must be met for approval.** To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- Member is 18 years of age or older and has a diagnosis of chronic kidney disease associated with type 2 diabetes
- Must submit lab test results documenting **BOTH** of the following obtained within the past 60 days
 - Members' current eGFR is > 25mL/minute/1.73 m²
 - Member's current Urinary Albumin-to -Creatinine Ratio (UACR) is ≥ 30 mg/g
- Member's current serum potassium is ≤5 mEq/L along with **BOTH** of the following (**submit current lab documentation obtained within the past 60 days**):
 - Therapy will **NOT** be initiated if serum potassium >5 mEq/L
 - Initiation with increased serum potassium monitoring during the first 4 weeks will be preformed if serum potassium is >4.8 to 5 mEq/L
- Member is established on treatment with maximally tolerated dose of angiotensin-converting enzyme inhibitor (ACE) or angiotensin receptor blocker (ARB) medication and will continue to take along with Kerendia® (finerenone)
- Member tried and failed, has a contradiction, or intolerance to at least **ONE** sodium glucose transport protein 2 (SGLT2) inhibitor that is indicated for use in patients with chronic kidney disease (e.g., Farxiga®, Jardiance®)
- Member does **NOT** have a diagnosis of adrenal insufficiency or a diagnosis of know significant non-diabetic renal disease, including clinically relevant renal artery stenosis
- Member is **NOT** receiving simultaneous treatment with strong CYP3A4 inhibitors
- For initial therapy, member will be dosed as follows:
 - eGFR ≥60 mL/minute/1.73 m²: starting dose will be 20 mg once daily
 - eGFR ≥25 to <60 mL/minute/1.73 m²: starting dose will be 10 mg once daily

(Continued on next page)

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member's current eGFR is >25 mL/minute/1.73 m² (**submit current lab documentation**)
- Member's Urinary Albumin-to-creatinine Ratio (UACR) has decreased by $\geq 30\%$ from baseline level and/or been sustained at $\geq 30\%$ below baseline level since last approval (**submit current lab documentation**)
- Member's current serum potassium level does **NOT** exceed 5.5 mEq/L (**submit current lab documentation**)
- Provider attests Kerendia[®] will be withheld if serum potassium is >5.5 mEq/L and will consider restarting therapy when serum potassium normalizes (≤ 5.0 mEq/L)

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****