

AvMed

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-877-535-1391. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If information provided is not complete, correct, or legible, authorization can be delayed.

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Drug Requested: Soliris® (eculizumab) IV (J1299) (Medical)
Atypical Hemolytic Uremic Syndrome (aHUS)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

- ☐ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

Recommended Dosage:

Patient Body Weight	Induction	Maintenance
40 kg and over	900 mg weekly for the first 4 weeks	1200 mg at week 5; then 1200 mg every 2 weeks
30 kg to less than 40 kg	600 mg for the first 2 weeks	900 mg at week 3; then 900 mg every 2 weeks
20 kg to less than 30 kg	600 mg for the first 2 weeks	600 mg at week 3; then 600 mg every 2 weeks
10 kg to less than 20 kg	600 mg single dose at week 1	300 mg at week 2; then 300 mg every 2 weeks
5 kg to less than 10 kg	300 mg single dose at week 1	300 mg at week 3; then 300 mg every 3 weeks

(Continued on next page)

Maximum Quantity Limit: 4 vials every 14 days; one 300 mg vial (30 mL) = 150 billable units [1 billable unit per 2 mg]

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- ☐ Prescribing physician must be or in consultation with a hematologist, oncologist, or nephrologist
- ☐ Prescriber must be enrolled in the Soliris® Risk Evaluation and Mitigation Strategy (REMS) program
- ☐ Member must be 2 months of age or older and has a weight of at least 5 kilograms
- ☐ Member must have a confirmed diagnosis of Atypical Hemolytic Uremic Syndrome (aHUS) (**must submit chart notes and labs**)
- ☐ Thrombotic Thrombocytopenic Purpura (TTP) has been ruled out by evaluating ADAMTS-13 level (ADAMTS-13 activity level >10%)
- ☐ Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS) has been ruled out
- ☐ Other causes have been ruled out such as coexisting diseases or conditions (e.g. bone marrow transplantation, solid organ transplantation, malignancy, autoimmune disorder, drug induced malignant hypertension, HIV infection, etc.) Streptococcus pneumonia or Influenza A (H1N1) infection, or cobalamin deficiency
- ☐ Documented baseline values of the following must be submitted: serum lactate dehydrogenase (LDH), serum creatinine/eGFR, platelet count, and plasma exchange/infusion requirement
- ☐ Member does **NOT** have a systemic infection
- ☐ Member must meet **ONE** of the following:
 - ☐ Member must be administered a meningococcal vaccine **at least two weeks prior** to initiation of Soliris® therapy and revaccinated according to current medical guidelines for vaccine use
 - ☐ Member has **NOT** received a meningococcal vaccination **at least two weeks prior** to the initiation of therapy with Soliris® and documented the risks of delaying Soliris® therapy outweigh the risks of developing a meningococcal infection
- ☐ Medication will **NOT** be used in combination with other complement inhibitor therapy (e.g., ravulizumab)

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member continues to meet all initial authorization criteria
- ☐ Member has **NOT** experienced unacceptable toxicity from the drug (e.g., serious meningococcal infections (septicemia and/or meningitis), infusion reactions, serious infections)

(Continued on next page)

- ☐ Provider must submit clinical notes **AND** labs documenting a positive clinical response or stabilization as evidenced by at least **ONE** of the following while on Soliris therapy (**check all that apply**):
- ☐ An increase in platelet count from baseline
 - ☐ Maintenance of normal platelet counts and LDH levels for at least 4 weeks
 - ☐ A 25% reduction in serum creatinine for a minimum of four weeks
 - ☐ Absence for at least 12 weeks of a decrease in platelet count of > 25% from baseline, plasma exchange or plasma infusion, and new dialysis requirement

EXCLUSIONS – Therapy will NOT be approved if member has history of any of the following:

- Unresolved meningococcal disease
- Any systemic bacterial or significant infections that have not been treated with appropriate antibiotics

Medication being provided by (check box below that applies):

- ☐ Location/site of drug administration: _____
NPI or DEA # of administering location: _____
- OR**
- ☐ Specialty Pharmacy

For urgent reviews: Practitioner should call AvMed Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. AvMed's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****