# AvMed

# PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

**Drug Requested: Pretomanid** 

MEM	BER & PRESCRIBER IN	<b>VFORMATION:</b> Authorization may be delayed if incomplete.
Member	Name:	
Member AvMed #:		Date of Birth:
Prescrib	er Name:	
		Date:
Office C	ontact Name:	
Phone Number:		Fax Number:
DEA OF	R NPI #:	
DRUG	GINFORMATION: Author	rization may be delayed if incomplete.
Drug Fo	rm/Strength:	
		Length of Therapy:
Diagnosis:		ICD Code, if applicable:
Weight:		Date:
support		below all that apply. All criteria must be met for approval. To tation, including lab results, diagnostics, and/or chart notes, must be
<u>Intial</u>	Authorization: 26 weeks	
□ T	he provider is an infectious dise	ease specialist or a pulmonologist
	AND	
n N	onresponsive multidrug-resistan Aycobacterium tuberculosis (Ple	lmonary extensively drug resistant (XDR), or treatment-intolerant, or at tuberculosis, NOT due to latent or extra-pulmonary infection due to ease submit chart note notes to include medical history and ics for detection of drug resistance)
	AND	
□ T	he patient had a chest x-ray cons	sistent with pulmonary tuberculosis (Please submit medical chart no
	AND	

amikacin) (Please submit pertinent medication history and medical chart notes)
pyrazinamide, ethambutol, a fluoroquinolone (such as levofloxacin) AND an injectable (such as
The patient's condition has been non-responsive to isoniazid, rifamycins (such as rifampin),

## **AND**

The patient been non-responsive to the best available regimen for at least 6 months

# <u>OR</u>

☐ The patient is intolerant or a contraindication with any of the following: para-amino salicylic acid, ethionamide, aminoglycosides (such as amikacin), or fluoroquinolones (such as levofloxacin)

#### **AND**

☐ Pretomanid will be taken in combination with bedaquiline (Sirturo®) and linezolid (Zyvox®) as part of the recommended dosing regimen, and will be administered by directly observed therapy (DOT)

## **AND**

☐ Prior to initiating combination therapy, the provider will monitor pertinent laboratory measures and assess for signs of liver injury, myelosuppression, and QT prolongation

<u>Reauthorization Approval</u>: 26 Additional Weeks. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

☐ The patient's infection of Mycobacterium tuberculosis requires further treatment (i.e. culture negative status was not observed at 6 months) (Please submit medical chart notes, culture results after initial 6 months treatment)

# <u>AND</u>

☐ Pretomanid will be taken in combination with bedaquiline (Sirturo®) and linezolid (Zyvox®) as part of the recommended dosing regimen - unless linezolid was discontinued after the first 4 weeks of consecutive treatment, then bedaquiline and pretomanid must be continued concomitantly

# **AND**

☐ The provider will continue to monitor pertinent laboratory measures and assess for signs of liver injury, myelosuppression, and QT prolongation

# Medication being provided by Specialty Pharmacy - PropriumRx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*Previous therapies will be verified through pha rmacy paid claims or submitted chart notes. \*