AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: Isturisa® (osilodrostat)

ME	MBER & PRESCRIBER INFO	DRMATION: Authorization may be delayed if incomplete.
Meml	ber Name:	
Member AvMed #:		
Presci	riber Name:	
		Date:
Office	e Contact Name:	
	e Number:	
DEA (OR NPI #:	
	UG INFORMATION: Authorizat	
Drug	Form/Strength:	
Dosing Schedule:		Length of Therapy:
Diagn	osis:	ICD Code, if applicable:
Weigl	ht:	Date:
<u>Ouar</u>	ntity Limits: 180 tablets per 30 da	ys (10 mg tabs); 120 tablets per 30 days (1 mg & 5 mg tabs)
suppo		w all that apply. All criteria must be met for approval. To on, including lab results, diagnostics, and/or chart notes, must be
Initi	ial Authorization: 6 months	
	Member must be 18 years of age or o	lder
	Provider is an endocrinologist or neu	rosurgeon
	Member has a diagnosis of persistent	or recurrent Cushing's disease
	Member must meet at least ONE of the following (chart notes must be submitted to document diagnosis and surgical history or contraindication to surgery):	
	• •	surgery and must be at least 30 days post-surgery
	(conventional radiation) post-pitu	•
	☐ Member is contraindicated to sur	gery AND irradiation

(Continued on next page)

	Member must have failed <u>90 days</u> of therapy with <u>ONE</u> of the following medications (verified by chart notes or pharmacy paid claims):	
	□ ketoconazole	
	□ metyrapone	
	□ mitotane	
	Member must have current mean urine free cortisol levels (mUFC) > 3 times the upper limit of normal (ULN)	
	Member will <u>NOT</u> use concurrent Cushing's disease treatment with Isturisa [®] (e.g., ketoconazole, metyrapone, mifepristone, mitotane)	
	Member has been assessed for QTc prolongation/Torsade de Pointes, hepatic and renal impairment	
	Member is <u>NOT</u> taking glucocorticoids (e.g. prednisone, hydrocortisone)	
	For members with diabetes and/or hypertension, disease is adequately controlled	
	 Member does NOT have a history of any of the following: Congestive Heart Failure (CHF) Unstable angina Sustained ventricular tachycardia Clinically significant bradycardia Advanced heart block Acute myocardial infarction <1 year prior to starting Isturisa Clinically significant impairment in cardiovascular disease 	
ppc	uthorization: 12 months. Check below all that apply. All criteria must be met for approval. To ort each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be ded or request may be denied.	
	Member's current 24-hour urinary free cortisol level is below the upper limit of normal (labs must be submitted)	
	Improvements in quality of life have been maintained while on Isturisa® therapy	
	Member will continue to be monitored for QTc prolongation, hepatic and renal impairment	
[ed	ication being provided by Specialty Pharmacy - PropriumRy	

**Use of samples to initiate therapy does not meet step edit/preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *