AvMed

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-877-535-1391</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx. Additional indications may be covered at the discretion of the health plan.

<u>Drug Requested</u>: Nplate[®] (romiplostim) (J2802) (Medical)

MEMBER & PRESCRIBER INF	ORMATION: Authorization may be delayed if incomplete.
Member Name:	
Member AvMed #:	Date of Birth:
Prescriber Name:	
Prescriber Signature:	
Office Contact Name:	
Phone Number:	
NPI #:	
DRUG INFORMATION: Authoriz	cation may be delayed if incomplete.
Drug Form/Strength:	
Dosing Schedule:	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight (if applicable):	Date weight obtained:
	t, the timeframe does not jeopardize the life or health of the member mum function and would not subject the member to severe pain.

Dosing Limits:

- A. Quantity Limit (max daily dose) [NDC Unit]:
 - Nplate 125 mcg SDV for injection: 40 vials per 28 days
 - Nplate 250 mcg SDV for injection: 20 vials per 28 days
 - Nplate 500 mcg SDV for injection: 12 vials per 28 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- Immune (idiopathic) thrombocytopenia (ITP): 1250 billable units weekly
- Hematopoietic Syndrome of Acute Radiation Syndrome (HSARS): 1250 billable units x 1 dose
- Injection, romiplostim, 1 microgram; 1 billable unit = 1 mcg

Dosed for one-time administration at 10 mcg/kg subcutaneously

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months		
	Prescribed by a hematologist	
	Member is not on any other thrombopoietin receptor agonist or mimetic (e.g., Doptelet, Mulpleta, Promacta) or Tavalisse	
	Provider attests the requested medication will NOT be used as an attempt to normalize platelet counts	
	Platelet count has been drawn within the previous 28 days (please submit labs)	
	Applicable diagnosis criteria below has been completed	
	Diagnosis: Myelodysplastic Syndromes (MDS)	
	Member is at least 18 years of age	
	Member has lower risk disease [i.e., IPSS-R (Very Low, Low, Intermediate), IPSS (Low/Intermediate-1), WPSS (Very Low, Low, Intermediate)]	
	Member has severe or refractory thrombocytopenia (i.e., platelet count $\leq 20 \times 10^9/L$ or higher with a history of bleeding)	
	Member progressed or had no response to hypomethylating agents (e.g., azacitidine, decitabine), immunosuppressive therapy, or clinical trial	
	Diagnosis: Hematopoietic Syndrome of Acute Radiation Syndrome (HS-ARS)	
	Member has suspected or confirmed exposure to radiation levels > 2 gray (Gy)	

□ Diagnosis: Immune (idiopathic) thrombocytopenia (ITP)		
\square Member is at increased risk for bleeding as indicated by platelet count $< 30 \times 10^9 / L$		
□ FOR acute immune (idiopathic) thrombocytopenia (ITP)		
☐ Member is at least 18 years of age		
☐ Member has previously failed <u>ONE</u> of the following treatments for ITP:		
☐ Corticosteroids (prednisone 0.5-2.0 mg/kg/day, or dexamethasone 40 mg/day for 4 days)		
□ IVIG		
□ Splenectomy		
□ Other:		
□ FOR chronic ITP lasting at least 6 months		
☐ Member is 1 year of age or older		
☐ Member has previously failed <u>ONE</u> of the following treatments for ITP:		
 Corticosteroids (defined as not having a response to at least a 3-month trial or is corticosteroid-dependent) 		
□ IVIG		
□ Splenectomy		
□ FOR members currently established on therapy:		
□ Past medical history and laboratory documentation has been provided to show platelet level monitoring and procedural records – PROCEED TO RENEWAL AUTHORIZATION CRITERIA		
Reauthorization: 12 months. (All indications) Check below all that apply. All criteria must be me for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.		
□ Diagnosis: Myelodysplastic Syndromes (MDS)		
☐ Member is not experiencing unacceptable toxicity from the drug (e.g., thrombotic/thromboembolic complications, risk of progression of myelodysplastic syndromes to acute myelogenous leukemia)		
☐ Member has not developed acute myeloid leukemia (AML) (NOTE: Nplate induces an increase in immature white blood cells and peripheral blasts which is not indicative of development of AML)		
☐ Member has experienced disease response indicated by an increase in platelet count compared to pretreatment baseline (not to exceed 450 x 10 ⁹ /L), reduction in bleeding events, or reduction in platelet transfusion requirements		

□ Provider will adhere to the following dosage reduction recommendations (platelet count drawn with the previous 28 days must be submitted):

Adjust dose in 250 mcg increments (from 250 mcg every other week up to 1000 mcg weekly) based on platelet count response:

- Platelet count $< 50 \times 10^9$ /L for three consecutive weeks: Increase to the next highest dose level
- Platelet count > 450×10^9 /L: Withhold the dose, reinitiate at a reduced dose when platelet count is $<200 \times 10^9$ /L
- □ Diagnosis: Hematopoietic Syndrome of Acute Radiation Syndrome (HS-ARS)
 - ***** Coverage cannot be renewed
- □ Diagnosis: Immune (idiopathic) thrombocytopenia (ITP)
 - ☐ Member is not experiencing unacceptable toxicity from the drug (e.g., thrombotic/thromboembolic complications, risk of progression of myelodysplastic syndromes to acute myelogenous leukemia)
 - \square Member has experienced disease response indicated by the achievement and maintenance of a platelet count of at least 50×10^9 /L (not to exceed 400×10^9 /L) as necessary to reduce the risk for bleeding
 - □ Provider will adhere to the following dosage reduction recommendations (platelet count drawn with the previous 28 days must be submitted):

Adjust dose based on platelet count response:

- Platelet count $< 50 \times 10^9$ /L: Increase weekly dose by 1 mcg/kg.
- Platelet count >200 ×10⁹/L to ≤ 400 ×10⁹/L for two consecutive weeks: Reduce weekly dose by 1 mcg/kg.
- Platelet count > 400×10^9 /L: Withhold dose; assess platelet count weekly; when platelet count < 200×10^9 /L, resume with the weekly dose reduced by 1 mcg/kg.
- DISCONTINUE, if platelet count does not increase to a level sufficient to avoid clinically important bleeding after 4 weeks at the maximum recommended dose of 10 mcg/kg/week.

Medication being provided by (check box below that applies):		
Location/site of drug administration:		
NPI or DEA # of administering location:		
<u>OR</u>		

☐ Specialty Pharmacy – Proprium Rx

Nplate (Medical) (AvMed) (Continued from previous page)

For urgent reviews: Practitioner should call AvMed Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. AvMed's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *