## **AvMed**

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

**<u>Drug Requested</u>**: Camzyos<sup>®</sup> (mavacamten)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.		
Member Name:		
Member AvMed #:		
Prescriber Name:		
	Date:	
Office Contact Name:		
Phone Number:	Fax Number:	
DEA OR NPI #:		
DRUG INFORMATION: Author	rization may be delayed if incomplete.	
Drug Form/Strength:		
	Length of Therapy:	
Diagnosis:	ICD Code:	
Weight:	Date:	
Quantity limit: 1 capsule per day		
	elow all that apply. All criteria must be met for approval. To support cluding lab results, diagnostics, and/or chart notes, must be provided	
Initial Authorization: 8 months		
☐ Member is 18 years of age or olde	er	
☐ Prescribed by or in consultation with a cardiologist specialist		
☐ Member has a diagnosis of symptomatic obstructive hypertrophic cardiomyopathy (HCM)		
☐ Member had an adequate echocardiogram or cardiovascular magnetic resonance imaging (CMR)		
☐ Member has New York Heart Association (NYHA) class II-III symptoms		

(Continued on next page)

	Baseline peak oxygen consumption (pVO2) determined by cardiopulmonary exercise testing (CPET) has been submitted	
	Member has documented left ventricular ejection fraction (LVEF) $\geq 55\%$	
	Member has a left ventricular outflow track (LVOT) gradient of 50 mmHg or higher	
	Member remains symptomatic despite trial of, or intolerant to at least <u>TWO</u> of the following (verified be chart notes or pharmacy paid claims):	
	☐ Beta-blocker (e.g., metoprolol, carvedilol)	
	☐ Calcium channel blocker (e.g., verapamil, diltiazem)	
	□ disopyramide	
	□ Septal reduction therapy	
	Member will avoid concomitant use with moderate to strong CYP2C19 inhibitors/inducers, strong CYP3A4 inhibitors/inducers	
	Member will avoid concomitant dual therapy with beta-blockers and calcium channel blockers or monotherapy with disopyramide or ranolazine	
	Member will <b>NOT</b> take disopyramide in conjunction with the requested medication	
ach l	uthorization: 12 months. All criteria that apply must be checked for approval. To support line checked, all documentation (lab results, diagnostics, and/or chart notes) must be provided juest may be denied.	
	Member has experienced continued clinical benefit as demonstrated by at least <b>ONE</b> of the following:	
	□ Improvement of at least 1.5 mL/kg/min in peak oxygen consumption (pVO2) as determined by cardiopulmonary exercise testing (CPET) AND a reduction of ≥ 1 New York Heart Association (NYHA) functional classification (e.g., I, II, III, or IV)	
	☐ Improvement of at least 3.0 mL/kg/min in pVO2 with no worsening in NYHA functional classification	
	Member has $\underline{NOT}$ experienced any treatment-restricting adverse effects (e.g., heart failure, LVEF <50% while taking requested medication	
	Provider has submitted the results of member's most recent echocardiogram or cardiovascular magnetic resonance imaging obtained after starting the requested medication	
<b>1ed</b>	ication being provided by a Specialty Pharmacy - PropriumRx	

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\*

\*Previous therapies will be verified throu.gh pharmacy paid claims or submitted chart notes.\*

<sup>\*</sup>Approved by Pharmacy and Therapeutics Committee: 7/21/2022; 5/25/2023 REVISED/UPDATED: 8/7/2022; 10/4/2022;06/15/2023; 10/26/2023