

# AvMed

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

### **Drug Requested:** (select drug below)

☐ Nexletol™ (bempedoic acid)

☐ Nexlizet™ (bempedoic acid/ezetimibe)

### **MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member AvMed #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

### **DRUG INFORMATION:** Authorization may be delayed if incomplete

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

### **Initial Authorization: 6 months**

- ☐ Must be prescribed or in consultation with one of the following:
  - ☐ Cardiologist
  - ☐ Endocrinologist
  - ☐ Lipid Specialist

(Continued on next page)

**DIAGNOSIS: select one below:**

- ☐ Atherosclerotic Cardiovascular Disease
- ☐ Heterozygous Familial Hypercholesterolemia (HeFH)
- ☐ Other diagnosis: \_\_\_\_\_
- ☐ ICD-10 Code(s) plus description: \_\_\_\_\_

- ☐ **Atherosclerotic Cardiovascular Disease** – Select if the member has Atherosclerotic cardiovascular disease (ASCVD) confirmed by the following: **(Please note: Chart documentation is required to be submitted along with this request form.)**

- ☐ Acute coronary syndrome
- ☐ History of myocardial infarction
- ☐ Stable or unstable angina
- ☐ Coronary artery disease
- ☐ Coronary or other arterial revascularization (e.g., percutaneous coronary intervention (PCI), angioplasty, coronary stent procedure or coronary bypass graft (CABG) surgery)
- ☐ Stroke
- ☐ Transient ischemic attack
- ☐ Peripheral arterial disease presumed to be of atherosclerotic origin

**OR**

- ☐ **Heterozygous Familial Hypercholesterolemia (HeFH)** – Select if the member has Heterozygous familial hypercholesterolemia (HeFH) confirmed by the following: **(Please note: Chart documentation is required to be submitted along with this request form.)**

- ☐ Untreated/pre-treatment LDL cholesterol (LDL-C)  $\geq$  190mg/dL in an adult or  $\geq$  155mg/dL in a child less than 16 years of age

**AND (ONE OF THE FOLLOWING)**

- ☐ Family history of myocardial infarction in first-degree relative less than 60 years of age
- ☐ Family history of myocardial infarction in second-degree relative less than 50 years of age
- ☐ Family history of familial hypercholesterolemia in first-or second degree relative
- ☐ Submission of medical records (e.g., chart notes, laboratory values) documenting LDL-C  $>$  190mg/dL in first or second degree relative.

**OR (ONE OF THE FOLLOWING)**

(Continued on next page)

- ☐ Genetic confirmation of functional mutation in the LDL receptor, Apo-B, or PCSK9 gene adaptor protein 1 (i.e., LDLRAP1 or ARH)
- ☐ Tendinous xanthomata
- ☐ Arcus cornealis before age 45

**AND**

**☐ Please confirm ALL the following for ASCVD and/or HeFH: (Please note: Chart documentation is required to be submitted along with this request form.)**

- ☐ Member is on high-intensity statin therapy (i.e., atorvastatin 40-80mg daily, rosuvastatin 20-40mg daily) **AND** ezetimibe concomitantly (unless the addition of ezetimibe is contraindicated) for > 12 continuous weeks (**Pharmacy claims will be verified**)

**OR**

- ☐ Member is unable to tolerate high intensity statin therapy and is on maximally tolerated statin therapy **AND** ezetimibe concomitantly (unless the addition of ezetimibe is contraindicated) for > 12 continuous weeks (**Pharmacy claims will be verified**)

**Statin:** \_\_\_\_\_ **Strength:** \_\_\_\_\_ **Date started:** \_\_\_\_\_

**AND (ONE OF THE FOLLOWING)**

- ☐ LDL-C remains greater than or equal to 70 mg/dL with ASCVD
- ☐ LDL-C remains greater than or equal to 100 mg/dL without ASCVD

**\*\*Please document: the LDL levels below (Labs MUST be attached or authorization will be delayed)\*\***

**LDL baseline:** \_\_\_\_\_ **LDL post therapy:** \_\_\_\_\_

**OR**

- ☐ Member is unable to tolerate statin therapy as evidenced by **ONE** of the following intolerable and persistent symptoms with **TWO** different statins (i.e., trial of at least 14 days of each) (**documentation MUST be provided and claims will be verified**):
  - ☐ Myalgia (muscle symptoms without CK elevations) **OR**
  - ☐ Myositis (muscle symptoms with CK elevations < 10 times upper limit of normal)

**AND**

- ☐ Reinitiating of statin therapy at lower dose or reduced frequency of administration must have been attempted and failed (**documentation of statin intolerance MUST be provided**)

**\*\*Please document statin therapy below; pharmacy claims will be verified\*\***

**Statin:** \_\_\_\_\_ **Strength:** \_\_\_\_\_ **Date started:** \_\_\_\_\_

**Statin:** \_\_\_\_\_ **Strength:** \_\_\_\_\_ **Date started:** \_\_\_\_\_

**OR**

(Continued on next page)

- ☐ Member has a labeled contraindication to ALL statins as documented in medical records and/or has experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations > 10 times upper limit of normal (**documentation of labeled contraindication to ALL statins must be provided**)

**AND**

- ☐ Member has had a **90-Day** trial of a PCSK9 inhibitor (i.e., Repatha® or Praluent® - require prior authorization) and failed to reach LDL target goal (**documentation of PCSK9 inhibitor failure, including LDL labs after 90 days of therapy, MUST be provided**)

**OR**

- ☐ Member has had a life-threatening adverse reaction to a PCSK9 inhibitors (i.e., Repatha® or Praluent® – required prior authorization) (**documentation of life-threatening adverse reaction MUST be provided**)

**\*\*Please note: Concomitant therapy with PCSK9 inhibitors will not be approved\*\***

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Documentation of positive clinical response to therapy (e.g., reduction in LDL-C levels)

**AND**

- ☐ Member continues to receive other lipid-lowering therapy (e.g., statin, ezetimibe) at a maximally tolerated dose

**OR**

- ☐ Member has a documented inability to take other lipid-lowering therapy (e.g., statins, ezetimibe)

**\*\*Please document: the LDL levels below (Labs **MUST** be attached or authorization will be delayed)\*\***

LDL baseline: \_\_\_\_\_ LDL post-Nexletol/Nexlizet: \_\_\_\_\_

**Medication being provided by a Specialty Pharmacy – Proprium Rx**

*Not all drugs may be covered under every Plan*

*If a drug is non-formulary on a Plan, documentation of medical necessity will be required.*

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****