AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Sodium Oxybate Products

<u>Drug Requested</u>: (select **ONE** from below)

□ Lumryz [™] (sodium oxbyate) ER oral suspension	□ Sodium Oxybate oral solution
□ Xyrem [®] (sodium oxybate) IR oral solution	□ Xywav [®] (sodium oxybate) low sodium IR oral solution
MEMBER & PRESCRIBER INFORMATI	ON: Authorization may be delayed if incomplete.
Member Name:	
Member AvMed#:	
Prescriber Name:	
Prescriber Signature:	
Office Contact Name:	
	Fax Number:
DEA OR NPI #:	
DRUG INFORMATION: Authorization may b	e delayed if incomplete.
Drug Form/Strength:	
Dosing Schedule: Length of Therapy:	
	ICD Code:
Weight:	Date:
	e drug's distribution is limited and prescribers

- To guard against diversion and misuse, the drug's distribution is limited and prescribers
 <u>MUST</u> adhere to a risk management protocol, the Lumryz[™]/Xyrem[®]/Xywav[®] REMS
 Program.
- All diagnosis of narcolepsy and Idiopathic Hypersomnia must be in accordance with the third edition of the International Classification of Sleep Disorders (ICSD-3), which is a fully revised version of the American Academy of Sleep Medicine's manual of sleep disorders nosology, published in cooperation with international sleep societies and is the key reference work for the diagnosis of sleep disorders.

• Lumryz[™]/Xyrem[®]/Xywav[®] for narcolepsy with or without cataplexy will not be approved in conjunction with Sunosi[®] or Wakix[®]. AvMed considers the use of concomitant therapy with Lumryz[™]/Xyrem[®]/Xywav[®] and Sunosi[®] or Wakix[®] to be experimental and investigational. Safety and efficacy of these combinations has not been established and will not be permitted. In the event a member has an active Sunosi[®] or Wakix[®] authorization on file, all subsequent requests for Lumryz[™]/Xyrem[®]/Xywav[®] will NOT be approved.

Recommended Dosage for Xyrem® & Xywav®:

Patient Weight	Initial Dosage		Maximum Weekly Dosage Increase		Maximum Recommended Dosage	
	Take at Bedtime	Take 2.5 to 4 Hours Later	Take at Bedtime	Take 2.5 to 4 Hours Later	Take at Bedtime	Take 2.5 to 4 Hours Later
<20 kg*	There is insufficient information to provide specific dosing who weigh less than 20 kg			recommendati	ions for patients	
20 kg to < 30 kg	≤1 g	≤1 g	0.5 g	0.5 g	3 g	3 g
30 kg to < 45 kg	≤1.5 g	≤1.5 g	0.5 g	0.5 g	3.75 g	3.75 g
≥45 kg	≤2.25 g	≤2.25 g	0.75 g	0.75 g	4.5 g	4.5 g

Recommended Dosage for LumryzTM:

• Starting dosage is 4.5 grams (g) once per night administered orally. Increase the dosage by 1.5 g per night at weekly intervals to the recommended dosage range of 6 g to 9 g once per night orally. The dosage may be gradually titrated based on efficacy and tolerability. Doses higher than 9 g per night have not been studied and should not ordinarily be administered.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Member must meet ONE of the following age requirements:
☐ For sodium oxybate/Xyrem [®] /Xywav [®] requests: Member is at least 7 years old
☐ For Lumryz [™] requests: Member is 18 years of age or older
Member's current weight must be noted if < 18 years old:kg
The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines
Must be prescribed by or in consultation with a neurologist, psychiatrist or sleep specialist
Member is NOT receiving treatment with sedative hypnotics or other CNS depressants (verified by paid pharmacy claims)
Member is NOT using alcohol
Member does NOT have a history of drug abuse

<u>m</u>	DIAGNOSIS: Narcolepsy with Cataplexy – Check below all that apply. All criteria must be net for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.
	Provider is requesting ONE of the following: □ Lumryz [™] □ sodium oxybate □ Xyrem [®] □ Xywav [®]
	Member has a diagnosis of narcolepsy with cataplexy (Multiple Sleep Latency Test (MSLT) confirming diagnosis of narcolepsy and chart notes documenting the occurrence of more than one episode of cataplexy at baseline prior to treatment with requested medication must be submitted. If polysomnography required, please submit with MSLT)
	Provider has submitted the member's baseline Epworth Sleepiness Scale score (rating scale must be attached)
	Member must have a <u>2-month</u> trial and failure of <u>ONE</u> of the following anti-cataplectic therapies (verified by pharmacy paid claims; documentation of <u>intolerance or treatment failure</u> must be submitted, unless use is contraindicated (please attach clinical documentation citing contraindication)):
	□ SSRI (i.e., fluoxetine)
	☐ TCA (i.e., clomipramine, imipramine, desipramine or protriptyline)
	□ SNRI (i.e., venlafaxine or duloxetine)
be de	PIAGNOSIS: Excessive Daytime Sleepiness associated with Narcolepsy — Check elow all that apply. All criteria must be met for approval. To support each line checked, all ocumentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be enied.
	Provider is requesting <u>ONE</u> of the following: □ Lumryz [™] □ sodium oxybate □ Xyrem [®] □ Xywav [®]
	Member has a diagnosis of excessive daytime sleepiness associated with narcolepsy (MSLT confirming diagnosis of narcolepsy must be submitted. If polysomnography required, please submit with MSLT)
	Member must have tried and failed at least 30 days of therapy with modafinil or armodafinil (verified by pharmacy paid claims; documentation of <u>intolerance or treatment failure</u> must be submitted)
	Provider has submitted the member's baseline Epworth Sleepiness Scale score (rating scale must be attached)

<u>fo</u>	IAGNOSIS: Idiopathic Hypersomnia – Check below all that apply. <u>All criteria must be met rapproval</u> . To support each line checked, all documentation, including lab results, diagnostics, and/or art notes, must be provided or request may be denied.
)	Provider is requesting <u>ONE</u> of the following: □ sodium oxybate □ Xyrem [®] □ Xywav [®]
ם	Member is at least 18 years old
	Member does NOT have cataplexy
3	Member has a diagnosis of idiopathic hypersomnia confirmed by <u>ALL</u> of the following ICSD-3 criteria (MSLT confirming diagnosis and polysomnography required, please submit with MSLT):
	☐ Member has < 2 sleep-onset rapid eye movement periods (SOREMPs) on a MSLT performed according to standard techniques, or has no SOREMPs if the REM sleep latency on the preceding nocturnal polysomnogram (PSG) was ≤ 15 minutes
	☐ Member has the presence of at least <u>ONE</u> of the following:
	\square Mean sleep latency of ≤ 8 minutes
	□ Total 24-hour sleep time ≥ 660 minutes (typically 12 to 14 hours) on 24-hour polysomnography monitoring or by wrist actigraphy in association with a sleep log
	☐ Insufficient sleep syndrome has been ruled out
	☐ The hypersomnolence and/or MSLT findings are not better explained by another sleep disorder, other medical or psychiatric disorder, or use of drugs or medications
3	Provider has submitted the member's baseline Epworth Sleepiness Scale score (rating scale must be attached)
3	Member must have a <u>2-month</u> trial and failure of <u>ONE</u> of the following Alerting Agents: □ amphetamine-based stimulant
	□ methylphenidate-based stimulant
)	Member must have a <u>2-month</u> trial and failure of <u>ONE</u> of the following Wake-promoting Agents: □ armodafinil (generic Nuvigil) □ modafinil (generic Provigil)
Tc	or <u>ALL</u> Xywav Requests – Check below all that apply. <u>All criteria must be met for approval</u> . support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, ust be provided or request may be denied.
3	Member must have a documented diagnosis of heart failure, renal failure or hypertension, are currently on a sodium restricted diet and are taking medications to control applicable diagnosis (verified by pharmacy paid claims)

Member must have an unsuccessful 30-day trial of sodium oxybate or Xyrem® (failure is defined as an
increase from baseline in ESS score and no change in cataplexy events from baseline); documentation
of intolerance or treatment failure must be submitted, unless use is contraindicated (please attach
appropriate clinical documentation)

For ALL Lumryz Requests – Check below all that apply. All criteria must be met for approval.
To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes,
must be provided or request may be denied.

Member must have an unsuccessful 30-day trial of sodium oxybate or Xyrem [®] (failure is defined as an
increase from baseline in ESS score and no change in cataplexy events from baseline); documentation
of intolerance or treatment failure must be submitted, unless use is contraindicated (please attach
appropriate clinical documentation)

Medication being provided by a Specialty Pharmacy – Proprium Rx

**Use of samples to initiate therapy does not meet step edit/preauthorization criteria. **

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.