AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (including phone and fax $\#_s$) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: Korlym[®] (mifepristone 300mg)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name:	
Member AvMed #:	Date of Birth:
Prescriber Name:	
Prescriber Signature:	Date:
Office Contact Name:	
	Fax Number:
DEA OR NPI #:	
DRUG INFORMATION: Authoriza	
Drug Form/Strength:	
Dosing Schedule:	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight:	Date:
Diagnosis:	ICD Code, if applicable:

Quantity Limits: 120 tablets per 30 days

Recommended Dosage: Initiate therapy with 300mg/day and titrate dose every 2-4 weeks based on tolerability and symptom control, and daily dose will NOT exceed 20mg/kg/day, OR 1,200 mg once daily

CLINICAL CRITERIA: Check below all that apply. <u>All criteria must be met for approval.</u> To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- □ Member is 18 years of age or older
- **D** Prescribing physician is an endocrinologist

(Continued on next page)

- □ Member has a diagnosis of Endogenous Cushing's Syndrome, and satisfies <u>ONE</u> of the following:
 - Diagnosis of Type 2 Diabetes Mellitus
 - □ Glucose intolerance noted by <u>ONE</u> of the following (must submit documentation): oral glucose tolerance test or Hemoglobin A1c test (HbA1c)
- □ Past medical history confirms <u>ONE</u> of the following:
 - □ Member has undergone surgery and has not been curative
 - □ Member is not a candidate for surgery
- Documentation of clinical failure (unable to normalize cortisol levels for the treatment of Cushing's Syndrome) to ketoconazole tablets taken along with <u>ONE</u> of the following:
 - □ An additional steroidogenesis inhibitor such as Metopirone (metyrapone capsules) or mitotane tablets
 - □ A pituitary-directed therapy such as cabergoline or Signifor LAR (pasireotide)
- □ Documentation of clinical failure to control glucose levels with Metformin AND <u>TWO</u> (2) of the following treatments:
 - Glucagon-Like Peptide-1 (GLP-1) Receptor Agonist (e.g., Trulicity, Ozempic)
 - □ Insulin (e.g., Humalog, Lantus)
 - Dipeptidyl Peptidase 4 (DPP-4) Inhibitor (e.g., Januvia, Onglyza)
 - Member is not also taking/ will not take strong inhibitors of CYP2A medications (e.g., simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, tacrolimus)
- □ For females of reproductive potential: pregnancy has been excluded before initiation of treatment and plans for prevention are implemented during treatment and for one month after stopping

<u>Reauthorization Approval</u>: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

□ Positive disease response has been achieved, demonstrated by improved glycemic control (decreased hemoglobin A1c) (current labs must be submitted to document HbA1c)

Medication being provided by a Specialty Pharmacy - PropriumRx

Not all drugs may be covered under every Plan.

If a drug is non-formulary on a Plan, documentation of medical necessity will be required. **Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.**

<u>Previous therapies will be verified through pharmacy paid claims or submitted chart notes.</u>